



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

common chord of circles O and O_1 , passing through A , MC and NC , respective tangents in M and N , B point where OM and O_1N meet, the five points F, B, M, C, N , are on the same circle. That B, M, N, C are on the same circle is evident. Then $\angle AFM = \angle CMA$, $\angle AFN = \angle ANC$.

$\therefore \angle MFN = \angle CMA + \angle CNA = 180^\circ - \angle C$, which proves that F is on circle $BMNC$, and therefore the proposition.

This is in all its generality, for we can readily see that B is the transformed of K_1K_2 —line joining points where the tangents meet the respective directrices.




Fig. 2.

360. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

A circular segment, area A , revolves successively about the diameters (fixed) d, d' , intersecting at an angle θ . If v = volume about d , v' the volume about d' , then $v^2 + v'^2 - 2vv' \cos \theta$ is independent of the position of the segment.

Solution by S. LEFSEHETZ, East Pittsburg, Pa., and the PROPOSER.

Let P be the center of gravity of the segment; EF , its chord; O the center of the circle; and AB, CD , the diameters d, d' , respectively; $\angle AOC = \angle BOD = \theta$.

Draw PQ perpendicular to CD , PM perpendicular to AB , QS perpendicular to PM , and OR perpendicular to QS .

Let $PQ = a$, $PO = c$, $PM = b$.

Then $b = PS + SM = PS + OR = a \cos \theta + \sqrt{c^2 - a^2} \sin \theta$. $v = 2\pi Ab$, $v' = 2\pi Aa$.

$\therefore v^2 + v'^2 - 2vv' \cos \theta = 4\pi^2 A^2 (a^2 + b^2 - 2abc \cos \theta) = \Delta$.

$$\therefore \Delta = 4\pi^2 A^2 [a^2 + a^2 \cos^2 \theta + 2a \sin \theta \cos \theta \sqrt{c^2 - a^2} + (c^2 - a^2) \sin^2 \theta - 2a^2 \cos^2 \theta - 2a \sin \theta \cos \theta \sqrt{c^2 - a^2}].$$

Hence, $\Delta = 4\pi^2 A^2 c^2 \sin^2 \theta$.

Solved similarly by S. G. Barton, J. Scheffer, and A. H. Holmes.

CALCULUS.

Remark on 282, by F. H. SAFFORD, Ph. D., University of Pennsylvania.

The published solution of 282 is incomplete. It is valid for a *long* box, but with a *short* box, the outer corner, B , may not reach its maximum before the corner, A , in Dr. Zerr's figure *emerges* from the hall. See Fig. 2, page 186, October, 1907. In that figure, N is omitted but should be vertically *above* Q . S is the corner vertically *under* M . In line 2, page 187, omit b preceding the word, becomes.